

# APP Intracellular Domain Impairs Adult Neurogenesis in Transgenic Mice by Inducing Neuroinflammation

Kaushik Ghosal, Andrea Stathopoulos, Sanjay W. Pimplikar\*

Department of Neurosciences, Lerner Research Institute, Cleveland Clinic, Cleveland, Ohio, United States of America

## Abstract

**Background:** A devastating aspect of Alzheimer's disease (AD) is the progressive deterioration of memory due to neuronal loss. Amyloid precursor protein (APP) occupies a central position in AD and APP-derived amyloid- $\beta$  ( $A\beta$ ) peptides are thought to play a pivotal role in disease pathogenesis. Nonetheless, it is becoming clear that AD etiology is highly complex and that factors other than  $A\beta$  also contribute to AD pathogenesis. APP intracellular domain (AICD) is generated together with  $A\beta$  and we recently showed that AICD transgenic mice recapitulate pathological features of AD such as tau hyperphosphorylation, memory deficits and neurodegeneration without increasing the  $A\beta$  levels. Since impaired adult neurogenesis is shown to augment memory deficits in AD mouse models, here we examined the status of adult neurogenesis in AICD transgenic mice.

**Methodology/Principal Finding:** We previously generated transgenic mice co-expressing 59-residue long AICD fragment and its binding partner Fe65. Hippocampal progenitor cell proliferation was determined by BrdU incorporation at 1.5, 3 and 12 months of age. Only male transgenic and their respective wild type littermate control mice were used. We find age-dependent decrease in BrdU incorporation and doublecortin-positive cells in the dentate gyrus of AICD transgenic mice suggesting impaired adult neurogenesis. This deficit resulted from decreased proliferation and survival, whereas neuronal differentiation remained unaffected. Importantly, this impairment was independent of  $A\beta$  since APP-KO mice expressing AICD also exhibit reduced neurogenesis. The defects in adult neurogenesis are prevented by long-term treatment with the non-steroidal anti-inflammatory agents ibuprofen or naproxen suggesting that neuroinflammation is critically involved in impaired adult neurogenesis in AICD transgenic mice.

**Conclusion/Significance:** Since adult neurogenesis is crucial for spatial memory, which is particularly vulnerable in AD, these findings suggest that AICD can exacerbate memory defects in AD by impairing adult neurogenesis. Our findings further establish that AICD, in addition to  $A\beta$ , contributes to AD pathology and that neuroinflammation plays a much broader role in AD pathogenesis than previously thought.

**Citation:** Ghosal K, Stathopoulos A, Pimplikar SW (2010) APP Intracellular Domain Impairs Adult Neurogenesis in Transgenic Mice by Inducing Neuroinflammation. PLoS ONE 5(7): e11866. doi:10.1371/journal.pone.0011866

**Editor:** Maria A. Deli, Hungarian Academy of Sciences, Hungary

**Received:** May 26, 2010; **Accepted:** July 4, 2010; **Published:** July 30, 2010

**Copyright:** © 2010 Ghosal et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** This work was supported by National Institutes of Health Grant R01-AG026146 and funding from Alzheimer's Association and CART-Rotary Funds to SWP. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: pimplis@ccf.org

## Introduction

Alzheimer's disease (AD) is a progressive neurodegenerative disorder characterized by a gradual decline in memory and executive functions. Mounting evidence suggests Amyloid precursor protein (APP) and presenilins are central molecules in the pathophysiology of AD [1,2]. Aided by presenilins, APP undergoes constitutive proteolysis to produce multiple fragments including amyloid  $\beta$  ( $A\beta$ ) peptides, which form senile plaques. A vast number of studies have shown that  $A\beta$  plays a pivotal role in the pathophysiology of AD [2] and it is thought that  $A\beta$  peptides, in an as yet uncertain form, trigger a cascade of downstream deleterious events that eventually results in AD.  $A\beta$  peptides are shown to hyperphosphorylate tau [3], induce neuronal cell death *in vitro*, and cause memory deficits in mice [4]. They are also implicated in causing silent seizures [5] and in the impairment of adult neurogenesis in AD mouse models [6–10].

The process of neurogenesis occurs mostly during embryogenesis but also persists during adulthood in the subventricular zone and subgranular zone (SGZ) of the hippocampus. There is increasing evidence that adult hippocampal neurogenesis plays an important role in learning and memory [11]. Increased adult neurogenesis enhances hippocampal-dependent learning and memory [12] whereas decreased neurogenesis impairs it [13]. Also, drugs that increase hippocampal neurogenesis also increase learning and memory [14] and hippocampal neurogenesis is known to decline with aging [15]. Adult neurogenesis is important for maintaining spatial memory function [13,16], which is characteristically lost in AD patients. Thus, impaired neurogenesis is being recognized as a significant pathological feature of AD since it can further exacerbate the memory decline caused by neuronal loss.

Although,  $A\beta$  peptides have been known to play a pivotal role in AD pathology, it is becoming increasingly clear that not all aspects of AD can be accounted for by  $A\beta$  alone [17,18]. Findings from

such diverse lines of investigation as neuroimaging, clinical trials and preclinical observations suggest that non-A $\beta$  factors also contribute to memory deficits in AD. The cleavage of APP by presenilins, which generates A $\beta$ , also releases APP Intracellular domain (AICD) from the membrane. AICD modulates gene transcription *in vitro* [19,20], alters signaling pathways and exerts deleterious effects in tissue culture cells [21,22], which could cause AD-like features *in vivo*. Indeed, AICD expressing transgenic mice recapitulate such AD pathologies as activation of glycogen synthase kinase-3 $\beta$  (GSK3 $\beta$ ), phosphorylation and aggregation of Tau, deficits in working memory and abnormal neural activity and silent seizures [23–25]. Interestingly, these pathologies were observed without appreciable changes in APP metabolism or A $\beta$  generation [24]. Since AICD levels are elevated in human AD brains [24], these findings raise the possibility that AICD, in addition to A $\beta$ , significantly contributes to AD pathogenesis.

In this study, we report that AICD transgenic mice also exhibit impaired adult neurogenesis by reducing neuronal proliferation and survival. This impairment is observed even in the absence of endogenous APP demonstrating that AICD alone is capable of inducing the deleterious effects. Importantly, the defective adult neurogenesis is prevented by blocking neuroinflammatory changes indicating that neuroinflammation plays a much broader role in AD pathogenesis than previously thought.

## Results

### Adult hippocampal progenitor cell (HPC) proliferation is reduced in AICD transgenic mice

We measured HPC proliferation in the SGZ by injecting male AICD transgenic mice (line FeC $\gamma$ 25) at 1.5, 3 or 12 months of age daily with BrdU for 3 days. Mice were sacrificed 24 hours after the last injection, brains were harvested, fixed and free-floating sections were prepared. We used male mice in this study to avoid the confounding effects of estrus cycle and gonadotrophins on adult neurogenesis in female animals [26]. Brain slices were immunostained with anti-BrdU antibody and BrdU-positive (BrdU+) cells were quantified in the SGZ (defined as the region spanning the border of granule cell layer and hilus with two cell layers into the granule cell layer and hilus) of the entire rostro-caudal extent of the hippocampus. At 1.5 months, the numbers of BrdU+ cells in FeC $\gamma$ 25 mice were similar to that observed in non-transgenic control littermates (Fig. 1A,B). However, 3-month-old FeC $\gamma$ 25 mice exhibited reduced BrdU+ cells compared to wild-type animals (Fig. 1C). Since FeC $\gamma$ 25 transgenic mice co-express AICD and Fe65 (which binds and stabilizes AICD), we performed an analogous experiment on Fe27 transgenic mice, which express Fe65 alone [23]. Fe27 mice showed a similar number of BrdU+ cells as the non-transgenic controls (Fig. 1C, E), indicating that AICD is required to observe impaired HPC proliferation in 3-month-old mice. The deficits in HPC proliferation were also observed in 12-month-old AICD transgenic mice ( $p < 0.05$ ; Fig. 1D, E). Animals older than 12 months were not studied because at this stage even the wild-type mice showed a very low level of adult neurogenesis. Together, these data suggest that AICD impairs adult HPC proliferation *in vivo*.

### Decreased hippocampal neurogenesis in AICD transgenic mice

A majority of newborn cells in the SGZ of adult animals do not survive, and the surviving cells differentiate into neurons or glia [27]. To examine whether reduced HPC proliferation is also accompanied by reduced neurogenesis, we immunostained the brain sections with anti-doublecortin (DCX) antibody. DCX is a

microtubule binding protein that is expressed by immature neurons but not by glia. At 3 months of age, wild-type mice showed a large number of highly branched DCX+ cells in the SGZ, with branch tips extending into the molecular layer (Fig. 2A). By contrast, the staining was severely diminished in age-matched FeC $\gamma$ 25 mice and there was a decrease in the number of DCX+ cells (Fig. 2A, 2D) as well as in the extent of arborization (see Fig. 2C). As observed above, the reduction in the number of DCX+ persisted in 12-month-old mice (Fig. 2B, 2D). Collectively, these results demonstrate that reduced HPC proliferation is accompanied by decreased neurogenesis in AICD transgenic mice and that these deficits are apparent at 3 months of age and continue to at least 12 months of age.


### AICD impairs cell survival but not cell differentiation in hippocampus

We further corroborated the finding that AICD transgenic mice show impaired HPC proliferation by determining the cell survival and differentiation of the newborn cells. To measure cell survival, we injected mice with BrdU for three consecutive days and sacrificed them either one day or four weeks after the last injection (Fig. 3A). Most newborn granular cells die during this time period and the surviving cells differentiate into neurons or glia. Quantification of BrdU+ cells showed that survival of the newborn cells (the ratio of BrdU+ cells after 4 weeks to the BrdU+ cells after 1 day) was decreased ( $p < 0.01$ ) in FeC $\gamma$ 25 mice (Fig. 3B). Roughly 19% and 14% of newborn cells in wild-type animals and in FeC $\gamma$ 25 mice survived after 4 weeks respectively.

We also determined the differentiation of newborn cells into neurons and glia by performing double-immunostaining using neuronal and astrocytic markers (Fig. S1). Brain sections were incubated with anti-BrdU antibody and anti-NeuN antibody (to mark mature neurons) or anti-GFAP antibody to identify astrocytes. There were no significant differences in the percentage of 'NeuN+BrdU' double-labeled cells and approximately half of the newborn cells matured into neurons in both FeC $\gamma$ 25 and wild-type mice (Fig. 3C). Similarly, there were no significant differences in the proportion of BrdU+ cells that matured into astrocytes (Fig. 3D). Together, these results indicate that AICD decreases HPC proliferation and survival without affecting cell differentiation.

### AICD impairs HPC proliferation in mice lacking APP

Adult neurogenesis is reduced in many mouse models of AD and this reduction may be caused by multiple factors [reviewed in 6]. We previously showed that APP processing and A $\beta$  levels were unchanged in 3-month-old FeC $\gamma$ 25 mice compared to control mice [24]. Thus, A $\beta$  probably plays no role in reduced adult neurogenesis in FeC $\gamma$ 25 animals. However, to address this issue unambiguously, we crossed FeC $\gamma$ 25 mice with APP knockout animals and obtained AICD-expressing APP $^{-/-}$  mice. 3-month-old FeC $\gamma$ 25;APP $^{-/-}$  mice were injected with BrdU for 3 days, brains were harvested on the 4<sup>th</sup> day and sections were immunostained for BrdU (Fig. 4A, B) or DCX (Fig. 4C). The lack of APP did not seem to affect adult neurogenesis since the number of BrdU+ cells in APP $^{-/-}$  animals (Fig. 4B) was similar to that seen in wild-type animals (see Fig. 1B). However, there was a small but significant decline ( $p < 0.05$ ) in the number of BrdU+ cells in the SGZ of FeC $\gamma$ 25;APP $^{-/-}$  (Fig. 4A, B). These results unambiguously show that AICD, in the absence of endogenous A $\beta$ , can impair HPC proliferation *in vivo*. We examined the DCX+ immature neurons in the SGZ of these mice and found that FeC $\gamma$ 25;APP $^{-/-}$  animals displayed a reduced number of DCX+ cells compared to APP $^{-/-}$  mice at 3 months of age (Fig. 4C, 4D).




**Figure 1. AICD transgenic mice show decreased cell proliferation in the SGZ of hippocampus.** Representative images of BrdU immunostaining in the subgranular zone (SGZ) of the dentate gyrus performed on 1.5-month-old (A) 3-month-old (C) and 12-month old (D) wild-type and FeC $\gamma$ 25 animals one day after the final BrdU injection (100 mg/kg). (A–B) At 1.5 mo there was no statistically significant difference in the number of BrdU-labeled cells between wild-type and FeC $\gamma$ 25 mice, quantified in (B). (E) Quantitative analysis of the total number of BrdU+ cells throughout the entire rostro-caudal extent of the hippocampus in 3-month-old (top) and 12-month-old (bottom) animals reveal a statistically significant decrease in the number of BrdU+ cells in the SGZ of FeC $\gamma$ 25 mice compared to wild-type mice. (n=3, 4 and 4 for both wild-type and FeC $\gamma$ 25 mice at ages 1.5 mo, 3 mo and 12 mo respectively. n=3 for Fe27 mice at 3 mo. \* p<0.05, \*\* p<0.01). doi:10.1371/journal.pone.0011866.g001

Also, the DCX+ cells showed reduced arborization in FeC $\gamma$ 25;APP<sup>-/-</sup> animals (Fig. 4C, right panels).

We next determined the level of HPC proliferation in two established mouse models of AD. APPPS1 mouse model expresses ‘Swedish’ mutant APP and M146L mutant presenilin [28] whereas R1.40 mice express only ‘Swedish’ mutant APP [29]. We observed

a reduction in HPC proliferation in 3-month-old APPPS1 mice but not in R1.40 animals (Fig. S2) even though both mouse models show elevated levels of A $\beta$ . We do not know whether R1.40 mice show neurogenesis defects at older ages but at a minimum, these results show that an elevation in A $\beta$  alone does not necessarily result in reduced neurogenesis. In any case, our findings confirm



**Figure 2. FeC $\gamma$ 25 mice show a decreased number of immature neurons in the SGZ of dentate gyrus (DG).** **A–B**, Coronal sections from 3 mo (**A**) and 12 mo (**B**) brains of wild-type and FeC $\gamma$ 25 mice were immunostained for the immature neuronal marker doublecortin (DCX). There is an abundance of immature neurons in the SGZ of the granule cell layer in both the supra- and infra-pyramidal blades of the DG. Compared to the wild-type (top) there is a significant decrease in the number of DCX+ cells in the SGZ of FeC $\gamma$ 25 mice (bottom). **C**, Magnification of the boxed region in (**A**) show the diminished number of DCX+ cells in FeC $\gamma$ 25 mice at 3 mo of age with a concomitant decline in dendritic number. **(D)** Quantitative analysis of the total number of DCX+ cells throughout the entire rostro-caudal extent of the hippocampus in 3-month-old (left) and 12-month-old (right) animals reveal a statistically significant decrease in the number of DCX+ cells in the SGZ of FeC $\gamma$ 25 mice compared to wild-type mice. Scale bar = 100  $\mu$ m.  $n=4$  for all.


doi:10.1371/journal.pone.0011866.g002

the previous results of reduced adult neurogenesis in mouse AD models and extend them to show that AICD, without A $\beta$ , is also able to impair HPC proliferation.

#### Treatment with NSAIDs prevents the deficits in HPC proliferation and neurogenesis

We next explored the potential mechanisms underlying the inhibitory effects of AICD. Impairment of adult neurogenesis by AICD in FeC $\gamma$ 25 mice is probably not mediated by cell-

autonomous mechanisms because AICD expression is driven by the CaMKII $\alpha$  promoter, which is active only in mature neurons [30]. However, since a number of factors including environmental stress and neuroinflammation are known to inhibit adult neurogenesis [31,32] we determined whether FeC $\gamma$ 25 mice showed signs of neuroinflammation. Unexpectedly, we observed a dramatic increase in the recruitment of CD45+ microglia in 12 week-old FeC $\gamma$ 25 mice compared to wild-type mice (Fig. 5A). We also observed an increased expression of several proinflammatory



**Figure 3. AICD downregulates cell survival but does not affect differentiation of newly generated hippocampal cells.** **A**, Experimental design to measure HPC proliferation or HPC maturation. After the last BrdU injection, mice were sacrificed after 1 day (proliferation) or 1 month (maturation) and HPC survival was quantified as the ratio of the number of BrdU+ cells present at 4 weeks post-BrdU to the number of BrdU+ cells present one day after the last BrdU injection. **(B)** FeC $\gamma$ 25 mice (13.7, SEM 0.02) showed a significant reduction in the cell survival potential compared to wild-type mice (18.3, SEM 0.02) (\*  $p=0.012$ ,  $\chi^2$  test). **C–D**, Quantitative analysis of the number of BrdU+ cells that mature into neurons **(C)** and astrocytes **(D)** 4 weeks after birth. Analysis reveals that AICD does not influence the proportion of surviving cells that mature into neurons (NeuN+, wild-type = 54.8, SEM 0.04; FeC $\gamma$ 25 = 56.6, SEM 0.1) or astrocytes (GFAP+, wild-type = 14.0, SEM 0.02; FeC $\gamma$ 25 = 15.1, SEM 0.03).  $n=3$  for all. doi:10.1371/journal.pone.0011866.g003


cytokines raising the possibility that a proinflammatory environment in FeC $\gamma$ 25 mice was responsible for the reduction in adult neurogenesis. To test this hypothesis, we treated mice with ibuprofen, a non-steroidal anti-inflammatory drug (NSAID), by feeding animals a drug-containing diet (375 ppm) for 9 weeks beginning at 3 weeks of age. Figure 5A shows that this regimen of ibuprofen treatment completely blocked the recruitment and activation of microglia in 3-month-old mice. Concomitantly, the number of BrdU+ cells in ibuprofen-fed FeC $\gamma$ 25 mice increased significantly and were comparable to those seen in wild-type mice (Fig. 5B, C). Compared to FeC $\gamma$ 25 mice on the control diet, ibuprofen-fed FeC $\gamma$ 25 mice showed a 28% increase in the number of BrdU+ cells in SGZ ( $p<0.05$ ). The NSAID diet did not increase the number of BrdU+ cells in wild-type mice ( $p=0.31$ ). We also performed DCX immunostaining on ibuprofen-treated animals and noticed an increase in DCX+ cells in ibuprofen-treated FeC $\gamma$ 25 mice (Fig. 5D). After the ibuprofen treatment, the DCX+ cells in AICD transgenic mice were morphologically indistinguishable from those in wild-type mice (Fig. 5E).

Since ibuprofen is also known to possess off-target effects (non-COX2), we used another NSAID, naproxen, to see whether it was able to block neurogenesis defects in AICD mice. Like above, AICD mice were fed naproxen-containing diet for 9 weeks and the brains were analyzed (Figure 6 A–D). Indeed, treatment with naproxen also resulted in increased BrdU incorporation in the dentate gyrus (Figure 6A) and quantification showed that the deficits were completely rescued (Fig. 6B). Similarly, naproxen-treated mice showed increased DCX-staining that was comparable to that seen in wild-type mice (Figure 6C, D). Thus, these data establish that NSAID treatment rescues the deficits in adult neurogenesis in AICD mice by blocking neuroinflammation.


## Discussion

Here we show that AICD impairs adult hippocampal neurogenesis in an age-dependent manner. Adult neurogenesis was normal in 6-week-old AICD transgenic mice but was reduced at 3 months of age and the impairment continued until at least 12 months of age. This impairment was due to decreased cell proliferation and survival and not due to altered differentiation. The defective neurogenesis is mediated by non-cell-autonomous mechanisms involving proinflammatory changes. The important finding of this study is that impaired adult neurogenesis in AD is due to neuroinflammation and the novelty is that we provide compelling evidence that overexpression of AICD impairs adult neurogenesis without A $\beta$ .


An increasing number of studies indicate that adult neurogenesis plays an essential role in normal memory function [11,33]. Inhibition of adult neurogenesis impairs certain kinds of memory, such as spatial learning [13,16]. Since the loss of spatial memory is a prominent neurological feature of AD, we suggest that impaired adult neurogenesis is especially detrimental in AD. Indeed, a vast number of studies have demonstrated deficits in adult neurogenesis in amyloid-based mouse models of AD [6–10,31,34], although a conflicting study has also been reported [35]. In AICD mice, impaired neurogenesis is first observed at 3 months, which precedes memory deficits by about 4–5 months [24]. Interestingly, impaired adult neurogenesis is also detected much before the appearance of other AD-features in APP<sup>swE</sup>/PS1<sup>dE9</sup> [9] and 3xTgAD [14] mouse models of AD. Thus, deficient adult neurogenesis is a significant pathological factor in AD and our findings suggest that therapies against AD will be more effective when supplemented with treatments that stimulate adult neuro-



**Figure 4. The effect of AICD is independent of other APP metabolites including A $\beta$ .** **A–B**, Decreased cell proliferation in FeC $\gamma$ 25;APP<sup>-/-</sup> mice compared to APP<sup>-/-</sup> mice. Photomicrograph reveals a reduced number of BrdU+ cells in the SGZ of FeC $\gamma$ 25;APP<sup>-/-</sup> compared to APP<sup>-/-</sup> which is quantified in **(B)** throughout the entire rostro-caudal extent of hippocampus. **C**, There was also a decrease in the number of immature neurons (DCX+) in the FeC $\gamma$ 25;APP<sup>-/-</sup> mice compared to APP<sup>-/-</sup>. Insets are magnified on the right to reveal the decreased number of DCX+ cells and the resulting dendritic branches in comparable regions of the hippocampus. **(D)** Quantitative analysis of the total number of DCX+ cells throughout the entire rostro-caudal extent of the hippocampus in 3-month-old animals reveal a statistically significant decrease in the number of DCX+ cells in the SGZ of FeC $\gamma$ 25; APP<sup>-/-</sup> mice compared to APP<sup>-/-</sup> mice. \*  $p < 0.05$  (Student's T-test).  $n = 3$  for all. doi:10.1371/journal.pone.0011866.g004



**Figure 5. Non-steroidal anti-inflammatory drug (NSAID; ibuprofen) treatment rescued impaired adult hippocampal neurogenesis in FeC $\gamma$ 25 mice.** **A**, CD-45 immunohistochemistry showed that ibuprofen treatment for 2 months (right) can decrease neuroinflammation seen in FeC $\gamma$ 25 mice compared to wild-type (WT) mice (left) at 3 months of age. **B–C**, Ibuprofen treatment rescued decreased adult hippocampal cell proliferation in FeC $\gamma$ 25 mice at 3 months, quantified in **C**. **D–E**, The number of immature neurons detected by doublecortin immunoreactivity was also increased in FeC $\gamma$ 25 mice (bottom) compared to wild-type mice (top) after ibuprofen treatment. Comparable hippocampal regions are identified in box **D** and enlarged in **E** for wild-type (left) and FeC $\gamma$ 25 mice (right). Error bars represent S.E.M. Scale bar = 100  $\mu$ m. n=4 for untreated wild-type and FeC $\gamma$ 25 mice and n=3 and 4 for ibuprofen treated wild-type and FeC $\gamma$ 25 mice respectively. doi:10.1371/journal.pone.0011866.g005



**Figure 6. Naproxen treatment also rescued defective adult neurogenesis.** **A–B,** Naproxen treatment rescued decreased adult hippocampal cell proliferation in FeC $\gamma$ 25 mice at 3 months, quantified in **B**. **C–D,** The number of immature neurons detected by doublecortin immunoreactivity was also increased in FeC $\gamma$ 25 mice (bottom) compared to wild-type mice (top) after naproxen treatment. Comparable hippocampal regions are

identified in box **C** and enlarged in **D** for wild-type (left) and FeC $\gamma$ 25 mice (right). Error bars represent S.E.M. Scale bar = 100  $\mu$ m. n = 4 for WT, FeC $\gamma$ 25 and WT with naproxen and n = 6 for FeC $\gamma$ 25 with naproxen. **E**. A schematic of how amyloidogenic processing of APP could bring about deleterious effects via both AICD and A $\beta$  peptides. Amyloidogenic processing [1] of APP generates A $\beta$  and promotes AICD signaling [45]. A $\beta$  peptides impair adult neurogenesis possibly by causing [2] an imbalance in GABAergic and Glutamatergic transmissions [10] whereas AICD does so by inducing [3] neuroinflammation (this study). Multiple studies have also shown that both A $\beta$  and AICD peptide fragments cause [4] other AD-pathological (including tau hyperphosphorylation, neuronal cell loss, silent seizures, memory deficits) features in mice (dashed arrows). doi:10.1371/journal.pone.0011866.g006

genesis (such as exercise) or prevent impaired adult neurogenesis (such as anti-inflammatory drugs). Indeed, our preliminary studies show that NSAID treatment improves working memory deficits in the Y-maze paradigm in older (>14 month) AICD transgenic mice.

The finding that treatment with ibuprofen or naproxen prevented neurogenesis defects is of relevance to AD and supports the potential of NSAIDs as prophylactic therapeutic agents [36]. How AICD overexpression results in sustained neuroinflammation is currently under investigation but our preliminary findings suggest that AICD activates stress-kinase pathways and upregulates expression of proinflammatory cytokines. Increased levels of proinflammatory cytokine are reported in serum and CSF from AD patients [37,38] and neuroinflammation is recognized as a significant pathological event in AD [39,40]. A mutation in presenilin that alters APP processing inhibits adult neurogenesis *in vivo* by activating the microglia and elevating the cytokine levels [31]. Proinflammatory conditions are known to downregulate neurogenesis and blocking inflammation has been shown to restore impaired neurogenesis [32]. At present the reasons why proinflammatory environment impairs proliferation and survival more potently but leaves neuronal maturation more or less intact are not clear. Multiple studies have shown the inhibitory effects of individual inflammatory factors (such as TNF- $\alpha$ , IFN- $\gamma$ , IL-6) on proliferation but variable effects on maturation [reviewed in 41]. NPC proliferation is orchestrated by temporal and spatial cues and involves expression of various transcription factors. It seems likely that NPC proliferation is more sensitive to the gene expression changes brought about by proinflammatory cytokines. In any case, chronic neuroinflammation in AD is likely to impair adult neurogenesis and aggravate the loss of hippocampal neurons. Interestingly, retrospective epidemiological studies [36] have shown that long-term use of NSAIDs exert a protective effect against developing AD (although short-term prospective use of NSAID was not protective) and this effect is proposed to be mediated by modulation of A $\beta$  levels [42]. Since both drugs, ibuprofen and naproxen, protected against AICD-induced defects in adult neurogenesis, we propose that NSAIDs exert beneficial effects independent of A $\beta$  modulation. Although our data show that inflammation plays a pivotal role in impaired neurogenesis, it is possible that AICD impairs neurogenesis also by stimulating GSK-3 $\beta$  activity [23]. Activation of GSK-3 $\beta$  alters the wnt signaling pathway and perturbs neurogenesis in mice [43,44] and in AD patients [45].

The observation that AICD transgenic mice lacking A $\beta$  exhibit impaired neurogenesis further strengthens the evidence that AICD also contributes to AD pathology. Although a vast amount of data support the pivotal role of A $\beta$  peptides in AD pathogenesis [1,2], it has become clear that AD etiology is complex and that a single causative agent cannot account for all of the available data [17,18]. AICD causes apoptosis *in vitro* [21] and induces tau hyperphosphorylation and aggregation, aberrant neuronal activity, memory deficit and neurodegeneration *in vivo* [24,25]. Because AICD levels are also elevated in human AD brains [24], we had hypothesized that AICD makes significant contributions to AD pathogenesis, and the present study further supports this

hypothesis. Interestingly, a recent study showed that AICD signaling occurs predominantly through the amyloidogenic processing of APP [46]. This raises the possibility that the harmful effects of amyloidogenic processing are mediated by both increased A $\beta$  as well as elevated AICD (Fig. 6E) and may explain the disappointing results from A $\beta$ -focused clinical trials. Future studies are needed to assess the individual contribution of each peptide to AD pathogenesis.

In conclusion, our study shows that AICD transgenic mice exhibit impaired adult hippocampal neurogenesis in an A $\beta$ -independent manner. More importantly, NSAID treatment had a protective effect against the neurogenesis defects. Together, these findings provide evidence for non-amyloid mechanisms that potentially exacerbate neuronal loss in AD and suggest that inflammation plays a much broader role in AD pathogenesis than previously appreciated.

## Methods

### Ethics Statement

All experiments were approved by the Institutional Animal Care and Use Committee of The Cleveland Clinic (Protocol Number: 08355).

### Transgenic mice

Transgenic mice co-expressing the AICD-59 and Fe65 (FeC $\gamma$ 25 line) or Fe65 alone (Fe27) in C57BL/6 background have been described previously [23,24,47]. Both transgenes are active in the brain under the control of the CAMKII $\alpha$  promoter. For all experiments non-transgenic littermates or C57BL/6 mice were used as wild-type controls. APP $^{-/-}$  have been described in detail previously [48] APPS1 and R1.40 animals have been previously described in detail [28], [29]. To generate FeC $\gamma$ 25/APP $^{-/-}$  mice that would express AICD59 and Fe65 on APP null background, FeC $\gamma$ 25 hemizygous mice were bred with APP $^{-/-}$  mice to obtain FeC $\gamma$ 25/APP $^{+/-}$  mice in the F1 generation. Subsequently, FeC $\gamma$ 25/APP $^{+/-}$  mice were bred with APP $^{+/-}$  to produce F2 FeC $\gamma$ 25/APP $^{-/-}$ , FeC $\gamma$ 25/APP $^{-/+}$ , APP $^{-/-}$  and APP $^{+/-}$  mice. All animals were housed under standard conditions with 4–5 mice per cage. Table 1 shows the various genotypes of mice used with their treatments groups.

### 5-Bromo-2-deoxyuridine injection and tissue preparation

To avoid the effects of gender and gonadotropic hormones on adult neurogenesis, only male mice were used throughout the study. To determine cell proliferation, mice were injected once daily with 5-Bromo-2-deoxyuridine (BrdU; 100 mg/kg, i.p.) for three consecutive days. On the day following the last BrdU injection, animals were anesthetized and transcardially perfused with ice-cold phosphate-buffered saline (PBS) followed by PBS containing 4% paraformaldehyde. Brain were removed and fixed in PBS containing 4% paraformaldehyde overnight at 4°C, cryoprotected in 30% sucrose in PBS, embedded using OCT compound (Sakura Finetek, Torrance, CA, USA) and stored at -80°C. To determine long-term cell survival and maturation,

**Table 1.** Study Design.

Genotype	Age (month)	Treatment	Test (marker)	Animals (n)
Wild Type	1.5	Regular Chow	AHP Proliferation (BrdU) Immature Neuron (DCX)	3
		Regular Chow	AHP Proliferation (BrdU) Immature Neuron (DCX)	4
	3	Regular Chow	Survival/Maturation (BrdU/NeuN/GFAP)	3
		Ibuprofen	AHP Proliferation (BrdU) Immature Neuron (DCX)	3
		Naproxen	AHP Proliferation (BrdU) Immature Neuron (DCX)	4
		Regular Chow	AHP Proliferation (BrdU) Immature Neuron (DCX)	4
FeC $\gamma$ 25	1.5	Regular Chow	AHP Proliferation (BrdU) Immature Neuron (DCX)	3
		Regular Chow	AHP Proliferation (BrdU) Immature Neuron (DCX)	4
	3	Regular Chow	Survival/Maturation (BrdU/NeuN/GFAP)	3
		Ibuprofen	AHP Proliferation (BrdU) Immature Neuron (DCX)	4
		Naproxen	AHP Proliferation (BrdU) Immature Neuron (DCX)	6
		Regular Chow	AHP Proliferation (BrdU) Immature Neuron (DCX)	4
12	Regular Chow	AHP Proliferation (BrdU) Immature Neuron (DCX)	4	
Fe27	3	Regular Chow	AHP Proliferation (BrdU) Immature Neuron (DCX)	3
App $-/-$	3	Regular Chow	AHP Proliferation (BrdU) Immature Neuron (DCX)	3
FeC $\gamma$ 25;App $-/-$	3	Regular Chow	AHP Proliferation (BrdU) Immature Neuron (DCX)	3
APPPS1	3	Regular Chow	AHP Proliferation (BrdU)	3
R1.40	3	Regular Chow	AHP Proliferation (BrdU)	3

Age (months) represents the time when BrdU injection was carried out. All Ibuprofen and Naproxen treatments were for 9 weeks terminating at 3 months of age. n represents number of animals.

doi:10.1371/journal.pone.0011866.t001

animals were killed 28 days after the final injection of BrdU. The brains were processed as described above.

### Immunohistochemistry

Hemi-brains were fixed overnight in 4% paraformaldehyde in PBS, sunk in 30% sucrose and embedded in OCT. 30  $\mu$ m sagittal sections were cut and every 12<sup>th</sup> section was used. Antibodies used were BrdU (Abcam, 1:500) and doublecortin (SantaCruz, 1:1000). Remaining details are given in Supplemental Methods S1.

### Stereological counts

The total number of BrdU+ or DCX+ cells in the subgranular zone of the dentate gyrus was quantified using unbiased stereological methods. BrdU+ or DCX+ cells were counted from every sixth section using a 40X objective throughout the entire rostro-caudal extent of the dentate gyrus (bregma  $-1.0$  mm to  $-2.80$  mm). Cells were counted from both halves of the brain within the granular cell layer (GCL) and adjacent SGZ up to a two-cell body-wide zone along the border between the GCL and the hilus. The experimenter counting the cells was blinded to the genotype and experimental modification of the mice. The total number of cells was obtained by multiplying the number of BrdU+ or DCX+ cells with inter section interval and adding them together for the entire hippocampus.

### Non-steroidal anti-inflammatory drug (NSAID) diet

Ibuprofen or naproxen (both from Sigma Aldrich) was formulated into standard animal chow by Research Diets at a final concentration of 375 ppm. Male FeC $\gamma$ 25 and non-transgenic littermates were fed drug-supplemented or control chow for 9 weeks. Mice were killed at the end of the experimental period and processed for histology or biochemical analyses. For neurogenesis assays, animals were injected with BrdU for three consecutive days before sacrificing as described above.

### Statistics

Statistical analysis was performed using GraphPad prism 3.1. The results are expressed as mean  $\pm$  standard errors of mean (SEM).

Rest of the protocols is described in Supplemental Methods (Methods S1).

### Supporting Information

#### Methods S1 Supplemental Methods

Found at: doi:10.1371/journal.pone.0011866.s001 (0.03 MB DOC)

**Figure S1** Differentiation of newborn granular cells into neurons and astrocytes. Mice were injected with BrdU and sacrificed one month after the last injection. Brains were harvested and free-floating sections were incubated with anti-BrdU antibody with anti-NeuN antibody (A) or anti-GFAP antibody (B). Confocal image analysis was used to score the coexpression of BrdU (green) with the neuronal marker NeuN (A, red) or the astrocyte marker GFAP (B, red).

Found at: doi:10.1371/journal.pone.0011866.s002 (6.99 MB TIF)

**Figure S2** Decreased neurogenesis in other mouse models of AD. A, Thioflavin-S staining of 3-month-old animals showed increased plaque density in APPPS1 animals but a complete absence in FeC $\gamma$ 25 transgenic animals. B, BrdU immunostaining on 3-month-old animals showed decreased cell proliferation in APPPS1 mice similar to FeC $\gamma$ 25 mice. R1.40 animals do not show a decline and behaved similar to WT animals. C, Quantification of BrdU counts throughout the entire hippocampus revealed a significant decrease in cell proliferation for both FeC $\gamma$ 25 and APPPS1 animals compared to R1.40 and WT mice. \*  $p < 0.05$  ANOVA. N = 4 for WT and FeC $\gamma$ 25 and 3 for APPPS1 and R1.40. Scale bar = 100  $\mu$ m

Found at: doi:10.1371/journal.pone.0011866.s003 (3.53 MB TIF)

## Acknowledgments

We thank Dr. Chris Nelson for his helpful suggestions and Dr. Bruce Lamb for providing us with mouse brain samples.

## References

- Price DL, Tanzi RE, Borchelt DR, Sisodia SS (1998) Alzheimer's disease: genetic studies and transgenic models. *Annu Rev Genet* 32: 461–493.
- Hardy J, Selkoe DJ (2002) The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics. *Science* 297: 353–356.
- Terwel D, Muyliaert D, Dewachter I, Borghgraef P, Croes S, et al. (2008) Amyloid activates GSK-3beta to aggravate neuronal tauopathy in bigenic mice. *Am J Pathol* 172: 786–798.
- Shankar GM, Li S, Mehta TH, Garcia-Munoz A, Shepardson NE, et al. (2008) Amyloid-beta protein dimers isolated directly from Alzheimer's brains impair synaptic plasticity and memory. *Nat Med* 14: 837–842.
- Palop JJ, Chin J, Roberson ED, Wang J, Thwin MT, et al. (2007) Aberrant excitatory neuronal activity and compensatory remodeling of inhibitory hippocampal circuits in mouse models of Alzheimer's disease. *Neuron* 55: 697–711.
- Lazarov O, Marr RA (2009) Neurogenesis and Alzheimer's disease: At the crossroads. *Exp Neurol*.
- Haughey NJ, Nath A, Chan SL, Borchard AC, Rao MS, et al. (2002) Disruption of neurogenesis by amyloid beta-peptide, and perturbed neural progenitor cell homeostasis, in models of Alzheimer's disease. *J Neurochem* 83: 1509–1524.
- Rodriguez JJ, Jones VC, Tabuchi M, Allan SM, Knight EM, et al. (2008) Impaired adult neurogenesis in the dentate gyrus of a triple transgenic mouse model of Alzheimer's disease. *PLoS ONE* 3: e2935.
- Demars M, Hu YS, Gadadhar A, Lazarov O. Impaired neurogenesis is an early event in the etiology of familial Alzheimer's disease in transgenic mice. *J Neurosci Res*.
- Sun B, Halabisky B, Zhou Y, Palop JJ, Yu G, et al. (2009) Imbalance between GABAergic and Glutamatergic Transmission Impairs Adult Neurogenesis in an Animal Model of Alzheimer's Disease. *Cell Stem Cell* 5: 624–633.
- van Praag H, Schinder AF, Christie BR, Toni N, Palmer TD, et al. (2002) Functional neurogenesis in the adult hippocampus. *Nature* 415: 1030–1034.
- Shors TJ, Miesegaes G, Beylin A, Zhao M, Rydel T, et al. (2001) Neurogenesis in the adult is involved in the formation of trace memories. *Nature* 410: 372–376.
- Snyder JS, Hong NS, McDonald RJ, Wojtowicz JM (2005) A role for adult neurogenesis in spatial long-term memory. *Neuroscience* 130: 843–852.
- Wang JM, Singh C, Liu L, Irwin RW, Chen S, et al. Allopregnanolone reverses neurogenic and cognitive deficits in mouse model of Alzheimer's disease. *Proc Natl Acad Sci U S A* 107: 6498–6503.
- Klempin F, Kempermann G (2007) Adult hippocampal neurogenesis and aging. *Eur Arch Psychiatry Clin Neurosci* 257: 271–280.
- Zhang CL, Zou Y, He W, Gage FH, Evans RM (2008) A role for adult TLX-positive neural stem cells in learning and behaviour. *Nature* 451: 1004–1007.
- Small SA, Duff K (2008) Linking Abeta and tau in late-onset Alzheimer's disease: a dual pathway hypothesis. *Neuron* 60: 534–542.
- Pimplikar SW (2009) Reassessing the amyloid cascade hypothesis of Alzheimer's disease. *Int J Biochem Cell Biol* 41: 1261–1268.
- Cao X, Sudhof TC (2001) A transcriptionally [correction of transcriptively] active complex of APP with Fe65 and histone acetyltransferase Tip60. *Science* 293: 115–120.
- Gao Y, Pimplikar SW (2001) The gamma-secretase-cleaved C-terminal fragment of amyloid precursor protein mediates signaling to the nucleus. *Proc Natl Acad Sci U S A* 98: 14979–14984.
- Passer B, Pellegrini L, Russo C, Siegel RM, Lenardo MJ, et al. (2000) Generation of an apoptotic intracellular peptide by gamma-secretase cleavage of Alzheimer's amyloid beta protein precursor. *J Alzheimers Dis* 2: 289–301.
- Kim HS, Kim EM, Lee JP, Park CH, Kim S, et al. (2003) C-terminal fragments of amyloid precursor protein exert neurotoxicity by inducing glycogen synthase kinase-3beta expression. *FASEB J* 17: 1951–1953.
- Ryan KA, Pimplikar SW (2005) Activation of GSK-3 and phosphorylation of CRMP2 in transgenic mice expressing APP intracellular domain. *J Cell Biol* 171: 327–335.
- Ghosal K, Vogt DL, Liang M, Shen Y, Lamb BT, et al. (2009) Alzheimer's disease-like pathological features in transgenic mice expressing the APP intracellular domain. *Proc Natl Acad Sci U S A* 106: 18367–18372.
- Vogt DL, Thomas D, Galvan V, Bredesen DE, Lamb BT, et al. (2009) Abnormal neuronal networks and seizure susceptibility in mice overexpressing the APP intracellular domain. *Neurobiol Aging*.
- Cameron HA, Gould E (1994) Adult neurogenesis is regulated by adrenal steroids in the dentate gyrus. *Neuroscience* 61: 203–209.
- Kempermann G, Gast D, Kronenberg G, Yamaguchi M, Gage FH (2003) Early determination and long-term persistence of adult-generated new neurons in the hippocampus of mice. *Development* 130: 391–399.
- Radde R, Bolmont T, Kaeser SA, Coomaraswamy J, Lindau D, et al. (2006) Abeta42-driven cerebral amyloidosis in transgenic mice reveals early and robust pathology. *EMBO Rep* 7: 940–946.
- Lamb BT, Bardel KA, Kulnane LS, Anderson JJ, Holtz G, et al. (1999) Amyloid production and deposition in mutant amyloid precursor protein and presenilin-1 yeast artificial chromosome transgenic mice. *Nat Neurosci* 2: 695–697.
- Abel T, Nguyen PV, Barad M, Deuel TA, Kandel ER, et al. (1997) Genetic demonstration of a role for PKA in the late phase of LTP and in hippocampus-based long-term memory. *Cell* 88: 615–626.
- Choi SH, Veeraraghavalu K, Lazarov O, Marler S, Ransohoff RM, et al. (2008) Non-cell-autonomous effects of presenilin 1 variants on enrichment-mediated hippocampal progenitor cell proliferation and differentiation. *Neuron* 59: 568–580.
- Monje ML, Toda H, Palmer TD (2003) Inflammatory blockade restores adult hippocampal neurogenesis. *Science* 302: 1760–1765.
- Aimone JB, Wiles J, Gage FH (2006) Potential role for adult neurogenesis in the encoding of time in new memories. *Nat Neurosci* 9: 723–727.
- Donovan MH, Yazdani U, Norris RD, Games D, German DC, et al. (2006) Decreased adult hippocampal neurogenesis in the PDAPP mouse model of Alzheimer's disease. *J Comp Neurol* 495: 70–83.
- Jin K, Galvan V, Xie L, Mao XO, Gorostiza OF, et al. (2004) Enhanced neurogenesis in Alzheimer's disease transgenic (PDGF-APPsw,Ind) mice. *Proc Natl Acad Sci U S A* 101: 13363–13367.
- in 't Veld BA, Ruitenber A, Hofman A, Launer LJ, van Duijn CM, et al. (2001) Nonsteroidal antiinflammatory drugs and the risk of Alzheimer's disease. *N Engl J Med* 345: 1515–1521.
- Ray S, Britschgi M, Herbert C, Takeda-Uchimura Y, Boxer A, et al. (2007) Classification and prediction of clinical Alzheimer's diagnosis based on plasma signaling proteins. *Nat Med* 13: 1359–1362.
- Lucin KM, Wyss-Coray T (2009) Immune activation in brain aging and neurodegeneration: too much or too little? *Neuron* 64: 110–122.
- Taupin P (2008) Adult neurogenesis, neuroinflammation and therapeutic potential of adult neural stem cells. *Int J Med Sci* 5: 127–132.
- Glass CK, Saijo K, Winner B, Marchetto MC, Gage FH. Mechanisms underlying inflammation in neurodegeneration. *Cell* 140: 918–934.
- Whitney NP, Eidem TM, Peng H, Huang Y, Zheng JC. Inflammation mediates varying effects in neurogenesis: relevance to the pathogenesis of brain injury and neurodegenerative disorders. *J Neurochem* 108(6): 1343–59.
- Weggen S, Eriksen JL, Das P, Sagi SA, Wang R, et al. (2001) A subset of NSAIDs lower amyloidogenic Abeta42 independently of cyclooxygenase activity. *Nature* 414: 212–216.
- Kim WY, Wang X, Wu Y, Doble BW, Patel S, et al. (2009) GSK-3 is a master regulator of neural progenitor homeostasis. *Nat Neurosci* 12: 1390–1397.
- Zhao C, Deng W, Gage FH (2008) Mechanisms and functional implications of adult neurogenesis. *Cell* 132: 645–660.
- He P, Shen Y (2009) Interruption of beta-catenin signaling reduces neurogenesis in Alzheimer's disease. *J Neurosci* 29: 6545–6557.
- Goodger ZV, Rajendran L, Trutzel A, Kohli BM, Nitsch RM, et al. (2009) Nuclear signaling by the APP intracellular domain occurs predominantly through the amyloidogenic processing pathway. *J Cell Sci* 122: 3703–3714.
- Ghosal K, Pimplikar SW (2010) Aging and excitotoxic stress exacerbate neural circuit reorganization in amyloid precursor protein intracellular domain transgenic mice. *Neurobiol Aging*;doi:10.1016/j.neurobiolaging.2010.04.020.
- Zheng H, Jiang M, Trumbauer ME, Sirimathsinghji DJ, Hopkins R, et al. (1995) beta-Amyloid precursor protein-deficient mice show reactive gliosis and decreased locomotor activity. *Cell* 81: 525–531.

## Author Contributions

Conceived and designed the experiments: KG SWP. Performed the experiments: KG AS. Analyzed the data: KG. Wrote the paper: KG SWP.