

Gene targeting by homologous recombination in mouse zygotes mediated by zinc-finger nucleases

Melanie Meyer^{a,b}, Martin Hrabé de Angelis^{b,c}, Wolfgang Wurst^{a,b,d,1}, and Ralf Kühn^{a,b,1}

^aInstitute for Developmental Genetics, Helmholtz Center Munich, Munich, Germany; ^bInstitute of Experimental Genetics, Helmholtz Center Munich, Munich, Germany; ^cTechnical University Munich, Munich, Germany; and ^dDeutsches Zentrum für Neurodegenerative Erkrankungen e.V., Munich, Germany

Communicated by Pierre Chambon, Institut de Génétique et de Biologie Moléculaire et Cellulaire, Illkirch-Cedex, France, June 30, 2010 (received for review June 16, 2010)

Gene targeting by homologous recombination in embryonic stem cells is extensively used to generate specific mouse mutants. However, most mammalian species lack tools for targeted gene manipulation. Since double-strand breaks strongly increase the rate of homologous recombination at genomic loci, we explored whether gene targeting can be directly performed in zygotes by the use of zinc-finger nucleases. Here we report that gene targeting is achieved in 1.7–4.5% of murine one-cell embryos upon the coinjection of targeting vectors with zinc-finger nucleases, without preselection. These findings enable the manipulation of the mammalian germ line in a single step in zygotes, independent of ES cells.

Gene targeting by homologous recombination in ES cells is widely used to mutate and modify the mouse genome. Thereby the mouse is established as the most commonly used animal model system (1). With the exception of mice, most other mammalian species lack tools for targeted gene manipulation because functional ES cell lines are not established. Gene targeting in somatic cells combined with nuclear transfer has been used as an alternative for the engineering of livestock species, but represents an elaborate task (2, 3).

Zinc-finger nucleases (ZFN) link a DNA binding domain of the zinc-finger type to the nuclease domain of Fok I and enable the induction of double-strand breaks (DSBs) at preselected genomic sites (4). DSBs closed by the error-prone, nonhomologous end-joining (NHEJ) DNA repair pathway frequently exhibit nucleotide deletions and insertions at the cleavage site. This technology has been applied to introduce knockout mutations into the germ line of rats and zebrafish by the expression of ZFNs in early embryos that target coding sequences (5–8).

It has been initially shown by using the yeast homing endonuclease I-SceI that the induction of DSBs at genomic insertions of I-SceI recognition sites increases the rate of homologous recombination (HR) in mammalian cells by several orders of magnitude (9). Artificially designed ZFNs further increased the ability to generate site-specific double-strand breaks in endogenous genes, without the requirement to introduce artificial nuclease recognition sites. Following this principle, ZFNs have been used to achieve efficient homologous recombination of gene targeting vectors with various endogenous loci in cultured and primary mammalian cells (10–13).

Zygotes provide a logical substrate for the single-step modification of the mammalian germ line. Whereas pronuclear DNA injection is routinely used to generate transgenic mice, rats, and other mammals by random genomic integration of transgenes (14–16), gene targeting in zygotes is not established due to the low rate of spontaneous HR. Because the mouse is the best established mammalian genetic model system, we explored whether targeted mutants can be directly obtained through ZFN-assisted homologous recombination in zygotes, upon the coinjection of ZFN mRNAs and gene targeting vectors (Fig. 1).




Fig. 1. Principle of ZFN-assisted gene targeting in zygotes. Zygotes collected from wild-type mice are coinjected into the pronucleus and cytoplasm with DNA of a gene targeting vector and mRNAs for the expression of a pair of gene specific zinc-finger nucleases (ZFN1/2). HR of the targeting vector with the target site results in a knockout (KO) or knock-in (KI) allele. Manipulated zygotes are subsequently transferred into pseudopregnant females to recover mutant mice.

Results

We used a ZFN pair (ZFN^{Rosa}) that recognizes 18 nucleotides upstream and 12 nucleotides downstream of the XbaI site in the first intron of *Rosa26* (Fig. 2A), a locus that enables widespread transgene expression in vivo (17). To confirm the stimulatory power of ZFN^{Rosa}-induced homologous recombination, we first transfected murine ES cells with a hygromycin selectable gene targeting vector (pRosa26.11; Fig. 2A) with or without ZFN^{Rosa} expression plasmids. Hygromycin-resistant ES cell clones were expanded and analyzed by Southern blotting for their *Rosa26* genotype. Thirty-two ES cell clones transfected with pRosa26.11 alone exhibited only the 11.5-kb EcoRV wild-type band (Fig. 2B). The cotransfection of pRosa26.11 with ZFN^{Rosa} plasmids strongly enhanced recombination at *Rosa26*, such that 11 of 34 analyzed clones exhibited the predicted 4.5-kb EcoRV fragment derived from a targeted allele (Fig. 2C). One ES cell clone exhibited only the targeted band and likely represents the targeting of both *Rosa26* alleles.

Targeted Integration of a β -Galactosidase Reporter Gene into the *Rosa26* Locus. To insert a β -Galactosidase gene into *Rosa26* of one-cell embryos, we linked a 4.2-kb reporter cassette to homology regions flanking the target XbaI site (Fig. 3A). This targeting vector was coinjected with ZFN^{Rosa} mRNAs in a two-step

Author contributions: M.H.d.A., W.W., and R.K. designed research; M.M. performed research; M.H.d.A. contributed new reagents/analytic tools; M.M. and R.K. analyzed data; and W.W. and R.K. wrote the paper.

The authors declare no conflict of interest.

¹To whom correspondence should be addressed. E-mail: wurst@helmholtz-muenchen.de or ralf.kuehn@helmholtz-muenchen.de.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1009424107/-DCSupplemental.




Fig. 2. Zinc-finger nuclease-assisted targeting of the *Rosa26* locus in ES cells. (A) Targeting vector pRosa26.11 for insertion of a hygromycin/puromycin resistance cassette, into the *Rosa26* locus. The structure of the wild-type locus, including the ZFN^{Rosa} recognition sites that overlap with an intronic XbaI site (X), and of the recombined *Rosa26* allele are shown. The location of the *Rosa26* promoter (Pr), first exon, of the 5'-*Rosa* Southern blot probe and of EcoRV (E) sites and fragments are indicated. (B) Southern blot analysis of EcoRV digested genomic DNA of 32 hygromycin resistant ES cell clones transfected with pRosa26.11 using the *Rosa26* 5' probe. The wild-type *Rosa26* locus exhibits a 11.5-kb EcoRV fragment (WT). (C) Southern blot analysis of EcoRV digested genomic DNA of 34 hygromycin resistant ES cell clones transfected with pRosa26.11 and ZFN^{Rosa} expression vectors using the *Rosa26* 5' probe. Targeted integration of the resistance gene cassette is indicated by the presence of an additional 4.5-kb EcoRV fragment.

procedure into the male pronucleus and the cytoplasm of mouse zygotes (see *Materials and Methods*). Injected zygotes were transferred into pseudopregnant females, and the resulting pups were analyzed at E18 for their *Rosa26* genotype. Southern blot analysis revealed the frequent loss of the XbaI site within the ZFN^{Rosa} target sequence in one or both *Rosa26* alleles, as indicated by the presence of a predicted 9-kb fragment in addition to the 4.7-kb wild-type band (Fig. 3B). Among 58 pups derived from these injections, we found 13 XbaI negative genotypes (Fig. 3B and Fig. S1A), indicating that mutagenic NHEJ events were induced by ZFN^{Rosa} in 22% of one-cell embryos. Some of these pups (Fig. 3B and Fig. S1A) had an intermediate pattern showing a majority of XbaI negative and a minor proportion of wild-type alleles, indicating that the ZFNs' action occurred in these embryos after the first cell division. Sequence analysis of PCR-amplified target regions confirmed that the loss of the XbaI site is caused by the deletion of multiple nucleotides within the ZFN target region (Fig. 3C). The DNA of pup #22 (Fig. 3E) showing a 8.5-kb XbaI band (Fig. 3B), predicted for homologous recombination (Fig. 3A), was further characterized by Southern blotting. This analysis proved to be a bona fide homologous recombination event, as indicated by the presence of the predicted 3.6-kb EcoRV and 9.4-kb EcoRV/SwaI fragments for pup #22 (Fig. 3B and D), detected with external 5'- and 3'-hybridization probes, respectively. The complete integration of the reporter gene in pup #22 was confirmed with an internal β -Galactosidase-specific probe that detected the predicted 8.5-kb XbaI band (Fig. 3D). The analysis of liver sections for β -Galactosidase activity proved

the functionality and heterozygous state of the targeted allele in pup #22 (Fig. 3E). In addition, we verified the integrity of the recombined locus by sequence analysis of the *Rosa26*/reporter junction regions amplified by PCR from pup #22 (Fig. S2A and B). Among the 58 analyzed pups, we found two, #14 and #21, that exhibit unexpected XbaI fragments of 4.7–8.5 kb (Fig. 3B) and were shown by Southern blot (Fig. 3D, #14) and PCR analysis (Fig. S2C) to contain 3' sequences, but not the coding region of the reporter cassette (Fig. 3D), and represent incomplete or rearranged recombination products.

Targeted Integration of a Venus Reporter Gene into the *Rosa26* Locus.

In a second experiment we used a targeting vector that inserts a 1.1-kb *Venus* reporter gene into the *Rosa26* locus (Fig. 4A). DNA analysis of 22 pups derived from zygote coinjections of this vector with ZFN^{Rosa} mRNAs revealed the presence of one recombined allele, as shown by a 3.1-kb BamHI and a 5.6-kb XbaI fragment (Fig. 4B, fetus 6, Fig. 4D) detected with the 5'-*Rosa* probe and by a 3.9-kb BamHI band detected with a *Venus* probe (Fig. 4C). The *Venus* reporter cassette was amplified from the genomic DNA of pup #6 and was found by sequence analysis to be entirely intact (Fig. S3). The second *Rosa26* allele of pup #6 had lost the XbaI site within the ZFN^{Rosa} target sequence, as indicated by the presence of a 9-kb XbaI fragment and the absence of the wild-type 4.7-kb band (Fig. 4B), detected with the *Rosa26* 5' probe.

Analysis of Random Vector Integrations. Pronuclear DNA injection is routinely used for the production of transgenic mice harboring random vector integrations (14). Unexpectedly, using Southern blotting and hybridization probes for the β -Galactosidase and *Venus* reporter gene coding regions, we did not observe any random integration of targeting vectors among the 80 pups derived from both ZFN coinjection experiments (Figs. 3D and 4C and Fig. S1B and C).

Discussion

Compared to a previous report on homologous recombination in mouse zygotes that described a spontaneous recombination rate below 0.1% (18), our results demonstrate that gene targeting can be achieved in pronuclei of one-cell embryos at a higher frequency (1.7–4.5%) when assisted by ZFNs. By analysis of the XbaI site located within the ZFN^{Rosa} recognition sequence, we determined that DSBs, which lead to NHEJ-induced loss of nucleotides, occurred in injected zygotes at an even higher rate of 22%. It has been demonstrated in *Drosophila* embryos that the suppression of NHEJ DNA repair leads to the preferential resolution of ZFN-induced DSBs by HR with a gene targeting vector, resulting in the increased recovery of targeted alleles (19). It is therefore possible that also in mammalian zygotes ZFN-assisted gene targeting can be further improved by the transient inhibition of the NHEJ pathway.

In the only reported case of HR in zygotes, the targeted allele exhibited numerous point mutations that interfered with gene expression, possibly as a result of an error-prone HR process in the pronuclei (18). Here we did not observe mutations in the analyzed regions of both targeted alleles, indicating that ZFN-assisted HR can occur with high fidelity as known for gene targeting in ES cells. As judged by the comparison of Southern blot signals derived from wild-type and mutant *Rosa26* loci, both correctly targeted alleles appeared in the heterozygous state indicating that ZFN expression, DSB induction, and HR can occur in injected pronuclei within a few hours, before the completion of DNA replication and the fusion of both pronuclei. Among the 80 pups analyzed, we also observed three genotypes (#14, #15, and #21, Fig. 3B) that exhibit a contribution of mutant *Rosa26* alleles below 50%, indicating that a part of the HR and NHEJ repair events occur late, only in a single cell of a 2-cell or 4-cell embryo. Homozygously modified genotypes, which exhibit the loss of the




Fig. 4. Targeted integration of a Venus reporter gene into the *Rosa26* locus. (A) Targeting vector pRosa26.3 for insertion of a 1.1-kb Venus gene, including a splice acceptor (SA) and polyA site, into the *Rosa26* locus. The structure of the wild-type locus, including the ZFN^{Rosa} recognition sites that overlap with an intronic XbaI site (X), and of the recombined *Rosa26* allele are shown. The location of the *Rosa26* promoter (Pr.), first exon, of Southern blot probes and XbaI (X) and BamHI (B) sites and fragments are indicated. (B) Genomic DNA of E18 fetuses derived from zygotes coinjections of ZFN^{Rosa} and pRosa26.3 was digested with BamHI or XbaI and analyzed by Southern blotting using the *Rosa26* 5' probe. The wild-type *Rosa26* locus exhibits a 5.8-kb BamHI and a 4.7-kb XbaI band. Targeted integration of the reporter gene is indicated by the presence of a predicted 3.1-kb BamHI and a 5.6-kb XbaI fragment detected with the *Rosa26* 5' probe (fetus #6). The loss of the ZFN^{Rosa} target XbaI site by NHEJ in nonrecombined alleles results in a 9.0-kb band (fetus #6). (C) Analysis of BamHI digested DNA with the internal Venus probe shows a predicted 3.9-kb band from the recombined allele of fetus 6. (D) Habitus of fetus #6 (recombined) and #7 (wild type).

polyadenylated using the Poly(A) Tailing Kit (Ambion). The resulting mRNA was purified using the MEGAClear™ Kit (Ambion) before resuspension in injection buffer (10 mM Tris, 0.1 mM EDTA, pH 7.2). The mRNA concentrations were determined by absorption and gel analysis, and each ZFN mRNA was diluted in injection buffer to a working concentration of 2.5 ng/μL together with targeting vector and stored at -80°C.

Targeting vector pRosa26.8 (a derivative of pRosa26.5) (20) consists of the 1-kb 5' *Rosa26* homology arm and the 4-kb 3' homology arm (17), flanking an adenoviral splice acceptor sequence, followed by the 3.2-kb β-Galactosidase coding region and the 220-bp SV40 polyadenylation site from plasmid pCMVβ (Fig. 2A). Targeting vector pROSA26.3 is equal to pRosa26.8, except that it contains a 1.1-kb reporter cassette for expression of the Venus GFP protein. The homology arms of both targeting vectors were sequenced and found identical to the respective genomic sequence of the strain C57BL/6J. pRosa26.8 was linearized at the 5' end of the short homology arm with I-SceI, whereas pRosa26.3 was used circular. The DNAs were purified with the Qiaquick PCR Purification kit (Quiagen), precipitated, and dissolved in injection buffer at a working concentration of 5 ng/μL (pRosa26.8-2) or 15 ng/μL (pRosa26.3-3) together with ZFN^{Rosa} mRNA (2.5 ng/μL each) and stored at -80°C.

Mouse zygotes were obtained by superovulation of Friend virus B (FVB) females and matings to C57BL/6N males (Charles River). The next day zygotes were collected from oviducts and microinjected in M2 embryo medium following standard procedures (21) with a mixture of targeting vector and ZFN^{Rosa} mRNAs (2.5 ng/μL each) loaded into a single microinjection needle. For microinjection a two-step procedure was applied: A first aliquot of the DNA/RNA mixture was injected into the male pronucleus (to deliver the DNA vector, as used for the production of transgenic mice). Upon the withdrawal of the injection needle from the pronucleus, a second aliquot of the DNA/RNA mixture was injected into the cytoplasm to deliver the ZFN mRNA directly to the translation machinery. Injections were performed using a Leica micromanipulator and microscope and an Eppendorf FemtoJet injection device. Injected zygotes were transferred into pseudopregnant CD1 female mice and fetuses recovered at day E18 for further analysis. From the transfer

of 469 zygotes injected with targeting vector pRosa26.8 we obtained 64 fetuses (14% recovery), from which 58 were further analyzed by Southern blotting. For pRosa26.3 we obtained 22 fetuses from 120 transferred zygotes (18% recovery). Recovered fetuses showed normal development and were viable (Figs. 3E and 4D). Mice were handled according to institutional guidelines and housed in standard cages in a specific pathogen-free facility on a 12 h light/dark cycle with ad libitum access to food and water.

X-Gal Staining. One longitudinal half of E18 fetuses were fixed and stained with X-Gal as described in ref. 21.

Preparation of Genomic DNA. A segment of the other embryo half was used for the isolation of genomic DNA. For this purpose frozen tissue segments were homogenized in lysis buffer (50 mM Tris-HCl, pH 8.0; 100 mM EDTA, pH 8.0; 1% SDS; 100 mM NaCl; 350 μg/μL Proteinase K) and incubated at 55°C for 2 h under vigorous shaking. Upon tissue lysis phenol/chloroform (1:1) was added, and the mixture was vortexed and centrifuged for 10 min at 16,000 × g. The supernatant was transferred into new tubes and extracted with chloroform. The supernatant was mixed with 0.7 volumes isopropanol to precipitate the genomic DNA. Precipitated DNA was collected by centrifugation, and the pellet was washed with 70% ethanol. The pellet was air-dried, resuspended in 10 mM Tris/1 mM EDTA buffer (pH 7.5), and incubated overnight at room temperature.

Southern Blot Analysis. Genomic DNA (6 μg) was digested overnight with 30 units restriction enzyme in a volume of 30 μL and then redigested with 10 units enzyme for 2–3 h. Samples were loaded on 0.8% agarose gels in Tris (89 mM)/borate (89 mM)/EDTA (2 mM), pH 8.0 buffer and run at 55 V overnight. The gels were then denatured for 1 h in 1.5 M NaCl; 0.5 M NaOH, neutralized for 1 h in 0.1 M Tris-HCl pH 7.5; 0.5 M NaCl, washed with 2× SSC, and blotted for 48 h with 20× SSC on Hybond N⁺ membranes (GE Healthcare). The membranes were then washed with 2× SSC, UV-cross-linked, and stored at -20°C. For hybridization the membranes were preincubated in Church buffer (1% BSA, 1 mM EDTA, 0.5 M phosphate buffer, 7% SDS) for 1 h at

65 °C under rotation. The Rosa26 5' probe was isolated as a 460-bp EcoRI fragment from plasmid pCRII-Rosa5' probe; the Rosa26 3' probe was isolated as 660-bp EcoRI fragment from the plasmid pCRII-Rosa3' probe, as described (17, 20). As β -Galactosidase probe we used the 1,250-bp SacII/EcoRV coding fragment from pCMV β and as Venus probe the venus coding region, isolated as a 730-bp BamHI/EcoRI fragment from pCS2-venus, was used. DNA fragments used as hybridization probes were heat denatured and labeled with P³² marked dCTP (Perkin Elmer) using the high-prime DNA labeling kit (Roche). Labeled probe DNA was purified on MicroSpin™ 5-200 HR columns (GE Healthcare), heat denatured, added to the hybridization buffer and membranes were rotated overnight at 65 °C. The washing buffer (2 \times SSC, 0.5% SDS) was prewarmed to 65 °C, and the membranes were washed three times (5 min, 30 min, and 15 min) at 65 °C under shaking. Next, the membranes were exposed at -80 °C to Biomax MS1 films and enhancing screens (Kodak) for 1–5 days until development. Photos of autoradiographs were taken with a digital camera on a transmitting light table, and segments were excised with the Adobe Photoshop software.

PCR and Sequence Analysis. To analyze the Rosa26 alleles recombined with pRosa26.8 we amplified DNA from fetuses 22, 21, and 14 (coinjecting with ZFN^{Rosa} and pRosa26.8) using the primer pairs Rosa 5HA (5'-aaagtcgctctg agttgttat-3') and SA1 (5'-cgatccccagctactggaaag-3') or SA2 (5'-ttgcttagc aggtctttc-3') to amplify the 5'-homology arm/splice acceptor junction and the primer pairs Rosa 3HA (5'-cacaccagggttagccttaagcc-3') and pA1 (5'-tcg gatcctctagagtcgagg-3') or pA2 (5'-aaaccacaactagaatgcagtg-3') to amplify the junction of the SV40 polyA/3'-homology arm. Amplification was performed using Herculase polymerase (Invitrogen) in 100- μ L reactions with 38 cycles of 95 °C -30 s; 59 °C -1 min; 72 °C -1 min. In embryos with recombination events PCR products have the expected sizes: Rosa 5HA/SA1: 346 bp; Rosa 5HA/SA2: 365 bp; Rosa 3HA/pA1: 467 bp; Rosa 3HA/pA2: 625 bp. PCR products from embryos exhibiting homologous recombination events were directly sequenced (Sequiseive) and compared to the pRosa26.8 vector sequence using the Vector NTI software (Invitrogen).

Genomic DNA from fetuses 9, 14, and 21, which lost the XbaI site in the ZFN target region as found by Southern blotting, and of wild-type controls were further analyzed for NHEJ-induced mutations by PCR amplification

using the primer pair Rosa 5HA and Rosa 3HA (wild-type PCR product: 252 bp). These PCR products were cloned using the StrataClone Blunt PCR Cloning Kit (Stratagene). Plasmid DNA was isolated from with the Qiaprep Spin Miniprep Kit (Qiagen) and analyzed by digestion with EcoRI to confirm the presence of vector inserts and digested with XbaI to analyze for the loss of the ZFN^{Rosa} target XbaI site. Selected clones were sequenced using the T3 forward primer (Sequiseive) and analyzed for mutations in comparison to the Rosa26 wild-type sequence using the Vector NTI software.

To analyze Rosa26 homologous recombined alleles with pRosa26.3, we amplified DNA from fetus 6 (coinjecting with ZFN^{Rosa} and pRosa26.3) using the primer pair Rosa 5HA (5'-aaagtcgctctgagttgttat-3') and Rosa 3HA (5'-cacaccagggttagccttaagcc-3'). The PCR product (5HA/3HA: 1,402 bp) was directly sequenced with a single 950-bp read (Sequiseive) and compared to the pRosa26.3 vector sequence using the Vector NTI software.

Culture and Transfection of ES Cells. JM8.A3 (22) ES cells were cultured and transfected as described for IDG3.2 ES cells (20) with 20- μ g linearized targeting vector pRosa26.11 [derived from pRosa26.10 (20)] and 5 μ g of a Ku70 subfragment expression vector, together with or without 15 μ g of circular ZFN^{Rosa} expression plasmids. Resistant colonies were selected with 125 U Hygromycin/mL culture medium, picked, and further expanded for DNA analysis. Recombinant ES clones were identified by Southern blotting, using EcoRV digestion and the Rosa26 5' probe, by the presence of a predicted 4.5-kb fragment in addition to the 11.5-kb wild-type band.

Note Added in Proof. While this article was in print we obtained also live adult heterozygous targeted mice from coinjections of ZFN^{Rosa} and pRosa26.3.

ACKNOWLEDGMENTS. We thank C. Arndt, R. Kneuttinger, A. Krause, A. Tasdemir, and S. Weidemann for excellent technical assistance, Sigma for ZFN plasmids, and J. Schick and R. Köster for reading the manuscript. This work was supported by the European Union within the European Conditional Mouse Mutagenesis Program (LSHG-CT-2005-018931, to W.W.) and by the German Ministry of Education and Research within the Disease genes to Protein pathways German genome research network (NGFN Plus) project (01GS0858, to W.W. and R.K.) of the NGFN-Plus program.

1. Capecchi MR (2005) Gene targeting in mice: Functional analysis of the mammalian genome for the twenty-first century. *Nat Rev Genet* 6:507–512.
2. Lai L, Prather RS (2003) Creating genetically modified pigs by using nuclear transfer. *Reprod Biol Endocrin* 1:82.
3. McCreath KJ, et al. (2000) Production of gene-targeted sheep by nuclear transfer from cultured somatic cells. *Nature* 405:1066–1069.
4. Porteus MH, Carroll D (2005) Gene targeting using zinc finger nucleases. *Nat Biotechnol* 23:967–973.
5. Doyon Y, et al. (2008) Heritable targeted gene disruption in zebrafish using designed zinc-finger nucleases. *Nat Biotechnol* 26:702–708.
6. Geurts AM, et al. (2009) Knockout rats via embryo microinjection of zinc-finger nucleases. *Science* 325:433.
7. Mashimo T, et al. (2010) Generation of knockout rats with X-linked severe combined immunodeficiency (X-SCID) using zinc-finger nucleases. *PLoS One* 5:e8870.
8. Meng X, Noyes MB, Zhu LJ, Lawson ND, Wolfe SA (2008) Targeted gene inactivation in zebrafish using engineered zinc-finger nucleases. *Nat Biotechnol* 26:695–701.
9. Rouet P, Smih F, Jasin M (1994) Expression of a site-specific endonuclease stimulates homologous recombination in mammalian cells. *Proc Natl Acad Sci USA* 91:6064–6068.
10. Hockemeyer D, et al. (2009) Efficient targeting of expressed and silent genes in human ESCs and iPSCs using zinc-finger nucleases. *Nat Biotechnol* 27:851–857.
11. Porteus MH, Baltimore D (2003) Chimeric nucleases stimulate gene targeting in human cells. *Science* 300:763.
12. Santiago Y, et al. (2008) Targeted gene knockout in mammalian cells by using engineered zinc-finger nucleases. *Proc Natl Acad Sci USA* 105:5809–5814.
13. Urnov FD, et al. (2005) Highly efficient endogenous human gene correction using designed zinc-finger nucleases. *Nature* 435:646–651.
14. Palmiter RD, Brinster RL (1985) Transgenic mice. *Cell* 41:343–345.
15. Pursel VG, Hammer RE, Bolt DJ, Palmiter RD, Brinster RL (1990) Integration, expression and germ-line transmission of growth-related genes in pigs. *J Reprod Fert* 51:77–87.
16. Rexroad CE, Jr, Hammer RE, Behringer RR, Palmiter RD, Brinster RL (1990) Insertion, expression and physiology of growth-regulating genes in ruminants. *J Reprod Fert* 51:119–124.
17. Soriano P (1999) Generalized lacZ expression with the ROSA26 Cre reporter strain. *Nat Genet* 21:70–71.
18. Brinster RL, et al. (1989) Targeted correction of a major histocompatibility class II E alpha gene by DNA microinjected into mouse eggs. *Proc Natl Acad Sci USA* 86:7087–7091.
19. Bozas A, Beumer KJ, Trautman JK, Carroll D (2009) Genetic analysis of zinc-finger nuclease-induced gene targeting in *Drosophila*. *Genetics* 182:641–651.
20. Hitz C, Wurst W, Kuhn R (2007) Conditional brain-specific knockdown of MAPK using Cre/loxP regulated RNA interference. *Nucleic Acids Res* 35:e90.
21. Nagy A, Gertsenstein M, Vintersten K, Behringer R (2003) *Manipulating the Mouse Embryo* (Cold Spring Harbor Lab Press, Cold Spring Harbor, NY), 3rd Ed.
22. Pettitt SJ, et al. (2009) Agouti C57BL/6N embryonic stem cells for mouse genetic resources. *Nat Methods* 6:493–495.